

**Elemental Composition of Plankton
In The Atlantic Ocean off the Coast of Lagos, Nigeria.**

***Akanmu, R. T. and Onyema, I. C.**

Department of Marine Sciences, University of Lagos, Akoka, Lagos, Nigeria.

Mail.: titipopoola@gmail.com

ARTICLE INFO

Article History

Received:26 /3/2019

Accepted:29/5/2019

Keywords:

Plankton, sea,
diatom, Iron,
SEM/EDS.

ABSTRACT

The determination of the elemental composition of plankton is important for interpretation of functional states of the aquatic ecosystems. The present study is focused on the elemental composition of the plankton assemblages from the Atlantic Ocean off the coast of Lagos, Nigeria using scanning electron microscope - SEM (VEGA3 TESCAN) coupled with energy dispersive x-ray spectrometry - EDS (Quantax 200, Bruker EDX) at an accelerating voltage of 5 - 20 kV. Surface plankton samples were collected by horizontal tows with a plankton net (55 μ m pore mesh size) at approximately 15 km away from the seashore along the entire Lagos shoreline (180km). The result from the SEM/EDS technique showed that the plankton is composed majorly of 13 elements (O, C, Si, Al, Cl, Fe, Mg, Na, K, S, Ti, Ca and P). The mineral particles of calcium (0.2%), potassium (1.3%), sodium (1.5%), chlorine (5.0%), magnesium (1.8%) and iron (3.3%) as well as organogenic elements of oxygen (36.9%), carbon (27.8%), sulphur (1%) and phosphorus (0.2%) were quantified.

INTRODUCTION

Phytoplankton are functional groups of prokaryotic and eukaryotic organisms (Lopes dos Santos *et al.*, 2017) that inhabit near the water surface where there is sufficient light to support photosynthesis. Zooplankton on the other hand is a collection of animal forms of drifting organisms in both fresh and marine waters (Onyema and Okedoyin, 2017). Therefore, both phytoplankton and zooplankton standing crop per time constitute the plankton assemblage. The ecology of planktonic organisms is influenced by the elemental characterization of some trace metals such as Fe, Cu, Zn, Cd and Hg (Nekhoroshkov *et al.*, 2014). The ability of plankton to concentrate trace elements from its environment is used to indicate the level of aquatic pollution in different regions of the World Ocean (McCormick and Cairns, 1994). Elemental accumulation as a dynamic process in plankton depends on the concentration of elements in the environment and different processes induced by the environmental nutritional stress, which are determined by the functional state of each organism (Sheue *et al.*, 2003). An earlier report of Leonova *et al.* (2013) showed that the plankton responds to the presence of heavy metals in aquatic environments, as well as upon that some metals inhibit and some stimulate the growth of organisms.

Goals and approaches to the study of the elemental concentration of plankton vary from one researcher to the other. For instance, Leonova *et al.*, (2013) estimated the concentration of a wide range of elements in the White Sea zooplankton, phytoplankton and zooplankton from freshwater Siberia lakes (Leonova *et al.*, 2006) and considered their accumulation capacity and involving into geochemical processes. The elemental content of the Black Sea plankton has now been studied for over 50 years (Vinogradova and Petkevich, 1967; Rozhanskaya, 1983; Saenko, 1992). Similarly, Nekhoroshkov *et al.*, (2014) documented the concentrations of 45 elements in the coastal phytoplankton communities of the Black Sea coastal area near Sevastopol, Ukraine while Yau *et al.*, (2016) investigated 16 cultured tropical microalgae species.

There is no known information on elemental identification and quantification of the plankton from Nigerian coastal waters hence, the aim of this study was to determine the elements that are embedded in the plankton of the Atlantic Ocean off the coast of Lagos, Nigeria. This will help to evaluate the nutritional properties of the plankton in this marine environment.

MATERIALS AND METHODS

Description of Study Site:

The Atlantic Ocean off the coast of Lagos borders Nigeria to the south (Fig. 1). It lies between Latitude $6^{\circ}15'52.9''\text{N}$ and Longitude $4^{\circ}05'49.4''\text{E}$ and falls within the Barrier Lagoon complex (200 km). It is a marine environment and salinity is a major factor in the growth of some organisms (Onyema and Akanmu, 2017). The study stations were approximately 15 km away from the seashore along the coastline between Badagry and Ibeju Lekki local government area in Lagos state, Nigeria. Ten sampling stations were chosen to cover the entire length of Lagos state in the ocean. The dominant ocean currents operating within the region include the Benguella, Guinea, Equatorial and the Equatorial countercurrents (Longhurst, 2006; Nwankwo and Onyema, 2003). The climate is the wet equatorial type influenced by nearness to the equator and in the Gulf of Guinea. The climate of the area as experienced in the Lagos metropolis is influenced by two air masses namely: the tropical maritime and tropical continental air masses.




Fig. 1 The Coast off Lagos, Nigeria showing the Sampling Stations

Collection of Plankton Samples:

Plankton samples were collected at the ten stations in April, 2016 with a 55 μ m mesh size standard plankton net towed horizontally from a motorized boat for 5 minutes at low speed (<4 knots). The filtered plankton were emptied into well-labelled 250 ml plastic containers with screw caps.

Analysis of Plankton Samples:

0.5kg concentrated plankton sample without acid washing step were fixed in 2% glutaraldehyde (C₅H₈O₂) that was prepared in 0.1 M sodium cacodylate buffer (C₂H₆AsNaO₂) at pH 7.2 for 48 hours at 4 °C. Samples were filtered on paper filters with 10 μ m pore size in the laboratory conditions. The sea-salt particles were dissolved and washed out after the treatment with distilled water. Then the filters with plankton were dried at room temperature. The samples were mounted on stubs using double-sticky tapes and then carbon (~10nm thick) coated at 20 mA for 180 s using sputter coater (BAL-TEC SCD005). Vacuum-evaporated carbon has a minimal influence on X-ray intensities on account of its low atomic number, and (unlike gold, which is commonly used for SEM specimens) does not add unwanted peaks to the X-ray spectrum. Samples were inserted into the working chamber for its elemental analysis at the reactor. Images of the samples were visualised and their elemental profiles were analysed using variable pressure scanning electron microscopy (model: VEGA3 TESCAN) at an accelerating voltage of 5 - 20 kV and a working distance of 7 - 15 mm. The microscope was coupled with energy dispersive X-ray spectrometry (Quantax 200, Bruker EDX) in order to obtain a distribution of elemental composition of the surface of plankton cell wall according to previously published data (Kania *et al.*, 2009; Kaliski *et al.*, 2012; Witek-Krowiak *et al.*, 2013; Yau *et al.*, 2016). The X-ray spectrum of the plankton loaded with a given microelement was obtained.

RESULTS

The plankton samples contain organic (phytoplankton, zooplankton and detritus) and inorganic particles. Therefore, scanning electron microscope (SEM/EDS) image showed aggregates of parts and tissues of plankton cells (Fig. 2) while diatoms of box and line shapes were revealed in SEM of plankton specimen (Fig. 3). A total of 13 elements in the order of oxygen (O), carbon (C), silicon (Si), aluminium (Al), chlorine (Cl), iron (Fe), magnesium (Mg), sodium (Na), potassium (K), sulphur (S), titanium (Ti), calcium (Ca) and phosphorus (P) were detected and distributed within the X-ray spectrum plankton (Fig. 4). In terms of percentage by weight of O, C, Si, Al, Cl, Fe, Mg, Na, K, S, Ti, Ca and P reported 37%, 28%, 14%, 7%, 5%, 3.3%, 1.8%, 1.5%, 1.3%, 1.0%, 0.3%, 0.2% and 0.2% respectively from the EDS. However, O, C, Si, Al, Cl and Fe were found to be higher than other elements comparatively (Fig. 5).




Fig. 2 The SEM image of the Morphology of the Scanned Sampled off the Coast of Lagos.




Fig. 3 SEM image of 10 μm plankton specimen off the coast of Lagos, Nigeria.




Fig. 4 X-ray Spectrum of the Elements in Plankton Specimen off the Coast of Lagos (EDS).




Fig. 5 Relative Levels of the Major Element Concentrations in the 100 µm Specimen off the Coast of Lagos, Nigeria (EDS).

DISCUSSION

The plankton from the Atlantic Ocean off the coast of Lagos was composed majorly of 13 elements oxygen (O), carbon (C), silicon (Si), aluminium (Al), chlorine (Cl), iron (Fe), magnesium (Mg), sodium (Na), potassium (K), sulphur (S), titanium (Ti), calcium (Ca) and phosphorus (P). Mineral or essential elements such as Ca, K, Na, Cl, Mg and Fe embedded in the plankton from this study have been reported to be responsible for the physiological processes of cells (White and Brown, 2010).

Previous studies of Yau *et al.*, (2016) detected 12 elements of Y, Nb, Fe, Ca, Cl, K, Cu, F, Ir, P, Mg and Si within 16 cultured tropical microalgae species using SEM/EDS while Nekhoroshkov *et al.*, (2014) reported concentration of 45 elements (Mg, Al, Cl, Ca, Ti, V, Mn, Cu, I, Na, K, Sc, Cr, Fe, Co, Ni, Zn, As, Se, Br, Rb, Sr, Zr, Mo, Ag, Sb, Cs, Ba, La, Ce, Nd, Sm, Gd, Tb, Tm, Yb, Hf, Ta, W, Au, Th, and U) determined by neutron activation analysis in plankton of the coastal zone of the Black Sea.

For this study, oxygen, carbon and silicon (78.6% by weight) were the most abundant element found in the sample. Organogenic elements such as O, C, S and P make up the total quantity or weight of living organisms. They form proteins, carbohydrates, lipids, nucleic acids and other biologically active compounds present in each cell of the phytoplankton (Nekhoroshkov *et al.*, 2014). Additionally, Priyadarshani and Rath (2012) reported that microalgae were rich in elements such as Fe, Ca, K and Mg and these additional nutrients made them suitable for nutraceutical products such as *Chlorella* sp. and *Spirulina* sp. Also, Ca, Mg, K, Cl and Na have been reported by (Nekhoroshkov *et al.*, 2014) to be the most abundant ions of seawater and by weight these ions constitute approximately 99% of sea salts while iron is the most important trace element that could increase phytoplankton production in ocean (Coale *et al.*, 1996). However, the high concentration of these elements in the environment

may slow down the vital processes, such as photosynthesis in the chloroplasts of phytoplankton cells (Nekhoroshkov *et al.*, 2014).

It is important to note that the high amounts of Si (13.9% by weight) recorded in this study could be attributed to the presence of diatoms within the phytoplankton spectrum. It has been reported by Yau *et al.* (2016) that silicon is an essential element to diatoms and it is required in the construction of their disc-like frustules. According to Onyema and Akanmu (2017) diatoms are the dominant phytoplankton groups in the sea off the coast of Badagry Lagos from microscopic study. Conclusively, mineral elements recorded for this study include Ca, Mg, K, Cl and Na are the most abundant ions found in seawater. The organogenic elements of O, C, S and P as well as Al, Si, Fe and Ti are nutrients required to sustain life in the ocean.

REFERENCES

- Coale, K. H., Johnson, K. S., Fitzwater, S. E., Gordon, R. M., Tanner, S., Chavez, F. P., Ferioli, L., Sakamoto, C., Rodgers, P., Millero, F., Steinberg, P., Nightingale, P., Cooper, D., Cochlan, W. P., Landry, M. R., Constantinou, J., Rollwagen, G., Trasvina, A. and Kudela, R. (1996). A massive phytoplankton bloom induced by an ecosystem-scale iron fertilization experiment in the equatorial Pacific Ocean. *Nature (Lond.)*, 383: 495 - 501.
- Kaliński, K., Marycz, K., Czogala, J., Serwa, E. and Janeczek, W. (2012). "An application of scanning electron microscopy combined with roentgen microanalysis (SEM-EDS) in canine urolithiasis," *Journal of Electron Microscopy*, 61: 47 – 55.
- Kania, M., Mikołajewska, D., Marycz, K. and Kobielarz, M. (2009). "Effect of diet on mechanical properties of horse's hair," *Acta of Bioengineering and Biomechanics*, 11(3): 53 – 57.
- Leonova, G. A., Bobrov, V. A. and Bogush, A. A. (2006). Estimation of the Geochemical Background and Anthropogenic Pollution of the Plankton by Microelement Composition (by the Example of the Western Siberia Lakes). *Bioindicators in Monitoring of Freshwater Ecosystems: Abstracts International Conference S-P.*, 91 - 92.
- Leonova, G. A., Bobrov, V. A., Bogush, A. A. and Bychinskii, V. A. (2013). Concentration of Chemical Elements by Zooplankton of the White Sea. *Marine Biology*, 53: 54 - 70.
- Longhurst, A. R. (2006). *Ecological Geography of the Sea*. (2nd ed.). Academic Press, San Diego, 123pp.
- Lopes dos Santos, A., Gourvil, P., Tragin, M, Noël, M. H., Decelle, J., Romac, S and Vaultot, D. (2017). Diversity and Oceanic Distribution of Prasinophytes Clade VII, The Dominant Group of Green Algae in Oceanic Waters. *The International Society of Microbial Ecology Journal*, 11: 512 - 528.
- McCormick, P. V. and Cairns, J. (1994). Algae as Indicators of Environmental Change. *Journal of Applied Phycology*, 6: 509 - 526.
- Nekhoroshkov, P. S., Kravtsova, A. V., Frontasyeva, M. V. and Tokarev, Y. N. (2014). Neutron Activation Analysis and Scanning Electron Microscopy of Phytoplankton in the Coastal Zone of Crimea (The Black Sea). *American Journal of Analytical Chemistry*, 5: 323 – 334.

- Nwankwo, D. I. and Onyema, I. C. (2003). A checklist of planktonic algae off Lagos coast. *Journal of Scientific Research Development*, 9: 75 – 82.
- Onyema, I. C. and Akanmu, R. T. (2017). Environmental Variables, Algal Pigments and Phytoplankton in the Atlantic Ocean Off the Coast of Badagry, Lagos. *Journal of Aquatic Sciences*, 32(1B): 171 - 191.
- Onyema, I. C. and Okedoyin, J. O. (2017). The Hydro-Environmental Conditions and Zooplankton of the Light House Creek, Lagos, Nigeria. *Journal of Aquatic Sciences*, 32(2): 285 - 301.
- Priyadarshani, I. and Rath, B. (2012). Commercial and industrial applications of micro algae – A review. *Journal of Algal Biomass Utilization*, 3(4): 89 - 100.
- Rozhanskaya, L. I. (1983). Trace Elements in Plankton and Fishes from the Western Part of the Black Sea. (inrussian). *Ekologiya Morya*, 12: 22 - 30.
- Saenko, G. N. (1992). Metals and Halogens in Marine Organisms. (inrussian). *Nauka. M. Acta*, 58: 123 - 134.
- Sheue-Duan, L., Pei-Chung, C. and Hoang-Kao, H. (2003) Benthic Algae as Monitors of Heavy Metals in Various Polluted Rivers by Energy Dispersive X-Ray Spectrometer. *Journal of Environmental Science and Health. Part A—Toxic/Hazardous Substances & Environmental Engineering*, 38A: 855-866.
- Vinogradova, Z. A. and Petkevich, T. A. (1967). Chemical Elemental Content of the Black Sea, Azov Sea and Caspian Sea Plankton. (inrussian). *Biochemistry of the Marine Organisms*, 8: 70 - 82.
- White, P. J. and Brown, P. H. (2010). Plant Nutrition for Sustainable Development and Global Health. *Annals of Botany*, 105: 1073 - 1080.
- Witek-Krowiak, A., Podstawczyk, D., Chojnacka, K., Dawiec, A. and Marycz, K. (2013). “Modelling and optimization of chromium (III) biosorption on soybean meal,” *Central European Journal of Chemistry*, 11(9): 1505 – 1517.
- Yau, S. K., Yusoff, F. M., Khong, N. M. H., Foo, S. C. and Lai, J. I. (2016). Improved pre-treatment protocol for scanning electron microscopy coupled with energy dispersive X-ray spectroscopy analysis of selected tropical microalgae. *Pertanika Journal of Tropical Agriculture Science*, 39(3): 321 – 342.